



# FARE.

Food Allergy Research & Education

## FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN

**PLACE  
PICTURE  
HERE**

Name: \_\_\_\_\_ D.O.B.: \_\_\_\_\_








Allergic to: \_\_\_\_\_

Weight: \_\_\_\_\_ lbs. Asthma:  **Yes (higher risk for a severe reaction)**  **No**

**NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.**

**Special Situation/ Circumstance - If this box is checked, the child has an extremely severe allergy to the following food(s)** \_\_\_\_\_  
**Even if the child has MILD symptoms after eating (ingesting) this food(s), Give Epinephrine immediately.**





For **ANY** of the following  
**SEVERE SYMPTOMS**

|   |  |   |   |
|---|--|---|---|
| <br><b>LUNG</b><br>Shortness of breath, wheezing, repetitive cough | <br><b>HEART</b><br>Pale or bluish skin, faintness, weak pulse, dizziness | <br><b>THROAT</b><br>Tight or hoarse throat, trouble breathing or swallowing     | <br><b>MOUTH</b><br>Significant swelling of the tongue or lips |
| <br><b>SKIN</b><br>Many hives over body, widespread redness        | <br><b>GUT</b><br>Repetitive vomiting, severe diarrhea                    | <br><b>OTHER</b><br>Feeling something bad is about to happen, anxiety, confusion | <b>OR A COMBINATION</b><br>of symptoms from different body areas  |

▼ ▼ ▼

1. **INJECT EPINEPHRINE IMMEDIATELY.**
2. **Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
- Consider giving additional medications following epinephrine:
  - » Antihistamine
  - » Inhaler (bronchodilator) if wheezing
- Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
- If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
- Alert emergency contacts.
- Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return

### MILD SYMPTOMS

|  |   |   |   |
|--|---|---|---|
| <br><b>NOSE</b><br>Itchy or runny nose, sneezing | <br><b>MOUTH</b><br>Itchy mouth | <br><b>SKIN</b><br>A few hives, mild itch | <br><b>GUT</b><br>Mild nausea or discomfort |
|--|---|---|---|

---

**FOR MILD SYMPTOMS FROM MORE THAN ONE BODY SYSTEM, GIVE EPINEPHRINE.**

---

**FOR MILD SYMPTOMS FROM A SINGLE BODY SYSTEM (E.G. SKIN, GI, ETC.), FOLLOW THE DIRECTIONS BELOW:**

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

### MEDICATIONS/DOSES

Epinephrine Brand or Generic: \_\_\_\_\_

Epinephrine Dose:  0.1 mg IM  0.15 mg IM  0.3 mg IM

Antihistamine Brand or Generic: \_\_\_\_\_

Antihistamine Dose: \_\_\_\_\_

Other (e.g., inhaler-bronchodilator if wheezing): \_\_\_\_\_

PATIENT OR PARENT/GUARDIAN AUTHORIZATION SIGNATURE \_\_\_\_\_ DATE \_\_\_\_\_ HEALTHCARE PROVIDER AUTHORIZATION SIGNATURE \_\_\_\_\_ DATE \_\_\_\_\_




# FARE.

Food Allergy Research & Education

## FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN


**HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO**

1. Remove Auvi-Q® from the outer case. Pull off red safety guard.
2. Place black end of Auvi-Q® against the middle of the outer thigh.
3. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
4. Call 911 and get emergency medical help right away.




**HOW TO USE EPIPEN®, EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN AUTO-INJECTOR, MYLAN**

1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, remove the blue safety release by pulling straight up.
3. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
4. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.




**HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENACLICK®), USP AUTO-INJECTOR, AMNEAL PHARMACEUTICALS**

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip. Grasp the auto-injector in your fist with the red tip pointing downward.
3. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh. Press down hard and hold firmly against the thigh for approximately 10 seconds.
4. Remove and massage the area for 10 seconds. Call 911 and get emergency medical help right away.




**HOW TO USE TEVA'S GENERIC EPIPEN® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR, TEVA PHARMACEUTICAL INDUSTRIES**

1. Quickly twist the yellow or green cap off of the auto-injector in the direction of the "twist arrow" to remove it.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, pull off the blue safety release.
3. Place the orange tip against the middle of the outer thigh at a right angle to the thigh.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



**HOW TO USE SYMJEPITM (EPINEPHRINE INJECTION, USP)**

1. When ready to inject, pull off cap to expose needle. Do not put finger on top of the device.
2. Hold SYMJEPITM by finger grips only and slowly insert the needle into the thigh. SYMJEPITM can be injected through clothing if necessary.
3. After needle is in thigh, push the plunger all the way down until it clicks and hold for 2 seconds.
4. Remove the syringe and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.
5. Once the injection has been administered, using one hand with fingers behind the needle slide safety guard over needle.



**ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:**

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

**OTHER DIRECTIONS/INFORMATION** (may self-carry epinephrine, may self-administer epinephrine, etc.):

Epinephrine first, then call 911. Monitor the patient and call their emergency contacts right away.

| EMERGENCY CONTACTS — CALL 911       | OTHER EMERGENCY CONTACTS              |
|-------------------------------------|---------------------------------------|
| RESCUE SQUAD: _____                 | NAME/RELATIONSHIP: _____ PHONE: _____ |
| DOCTOR: _____ PHONE: _____          | NAME/RELATIONSHIP: _____ PHONE: _____ |
| PARENT/GUARDIAN: _____ PHONE: _____ | NAME/RELATIONSHIP: _____ PHONE: _____ |